

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):


- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

Ataxin-2

- 120 kD cytoplasmic protein
- orthologs in various species



Body Weight






FIGURE 2

Body weight for the three genotypes with *ad lib* diet




Postnatal Age:
12.7 weeks


Background:

62.5%C57BL/6J/37.5%129X1/SvJ

Average Daily Food Intake for the three Genotypes under *ad lib* Conditions




Body weight for the three genotypes with restricted diet (4g per day)



Postnatal Age:
10.7 weeks

week

Average Daily Food Intake for the three Genotypes under restricted Conditions (4g/day)



Weight Gain for Male Mice:

Ad lib $-/-$ and $+/-$ mice are heavier than Wt

